

Total laparoscopic pancreaticoduodenectomy versus open pancreaticoduodenectomy (TJDBPS01): study protocol for a multicentre, randomized controlled clinical trial.

ABSTRACT

Introduction Pancreatoduodenectomy (PD) is one of the most complex abdominal operations to perform, and it is usually conducted for tumours of the periampullary region and chronic pancreatitis. Minimally invasive surgery has been progressively being developed for pancreatic surgery, first with the advent of hybrid-laparoscopy and recently with total laparoscopic surgery. Issues including the safety and efficacy of total laparoscopic pancreaticoduodenectomy (TLPD) and open pancreaticoduodenectomy (OPD) are currently being debated. Studies comparing these two surgical techniques are emerging, and large randomized controlled trials (RCTs) are lacking but are clearly required.

Methods and analysis TJDBPS01 is a multicentre, prospective, randomized controlled, parallel-group, superiority trial in fourteen centres with pancreatic surgery experts who have performed ≥ 104 TLPDs and OPDs. A total of 656 patients who will undergo PD are randomly allocated to the TLPD group or OPD group in a 1:1 ratio. The trial hypothesis is that TLPD has superior or equivalent safety and advantages in postoperative recovery compared with OPD. The primary outcome is the postoperative length of stay (LOS).

Ethics and dissemination The Institutional Review Board Approval of Tongji Hospital, Tongji Medical College, Huazhong University of Science and Technology has approved this trial and will be routinely monitoring the trial at frequent intervals, as will an independent third-party organization. Any results from this trial (publications, conference presentations) will be published in peer-

reviewed journals and conference proceedings.

Trial registration number NCT03138213 in clinicaltrials.gov.

Keywords pancreaticoduodenectomy, laparoscopic surgery, open surgery, whipple, randomized controlled trial.

Article Summary

Strengths and limitations of this study

- The study is designed as a prospective, randomized-controlled trial to achieve conclusion on the highest evidence level to provide the evidence concerning the possible benefits of total laparoscopic pancreaticoduodenectomy (TLPD) over open pancreaticoduodenectomy (OPD) in China, it is (i) multicentric and (ii) internationally registered.
- Only high volume, expert centres are qualified to participate the study. They have to provide (i) laparoscopically trained surgeons with ≥ 104 TLPD procedures performed, (ii) ≥ 104 OPD procedures performed and (iii) completed the Tongji Hospital TLPD training programme.
- This is the first randomized trial providing with a 2-year survival data after pancreaticoduodenectomy.
- The main limitation is as this study is carried out by a large team of physicians that consists of surgeons, gastroenterologists, radiologists, pathologists and oncologists in several districts, the coordination of the team becomes a big challenge.
- This trial is designed to compare two surgery techniques with an open label so that the patients and surgeons are not blinded for their interventions, however the data collectors and the analysts are blinded.

Word count 4999

INTRODUCTION

Pancreaticoduodenectomy (PD) is one of the most complex abdominal operations to perform, involving removal of the head of the pancreas, the duodenum and the common bile duct, and it is usually performed for the treatment of pancreatic cancer and the relief pain in chronic pancreatitis.^{1,2}

Introduced by Gagner et al, who performed the first total laparoscopic pancreaticoduodenectomy (TLPD) procedures in 1994, the use of laparoscopy in pancreas surgery has continued to grow rapidly because of its potential technical advantages, such as the precision of movements achieved using laparoscopic instruments and a three-dimensional view provided by the unique vision system.³

To date, many studies have confirmed the advantages of laparoscopic techniques in gastric, renal and cervical surgeries.⁴⁻⁶ However, only a few low-quality studies have reported the application of total laparoscopy in pancreas surgery or in PD surgery. Notably, recently performed meta-analyses comparing LPD and open PD (OPD) have been unsatisfactory due to the small number of studies of good quality.⁷⁻¹⁰ Most studies concluded that TLPD was associated with a reduction in estimated blood loss (EBL), a lower transfusion rate and a shorter length of stay (LOS) compared with OPD surgery. In addition, TLPDs have a much longer operative time than OPDs. Finally, there were no significant differences between the two procedures in terms of major complications or mortality.

Similarly, all of the studies that reported results on this topic emphasized the need for large randomized controlled trials (RCT). However, RCTs are difficult to perform and are costly. There should be further consideration regarding the need for detection of numerous surgical, clinical and

oncological variables. Thus, it is imperative for such a study to have enough patients enrolled. A study recruiting an adequate number of patients is desirable. In year 2017, Palanivelu et al published the first RCT worldwide and concluded that LPD offered a shorter hospital stay compared with OPD.¹¹ Poves reached the same conclusion the following year.¹² Both of their studies were conducted in a single-centre setting and recruited only 60 and 66 patients, respectively. The researchers claimed that a multicentre trial will be necessary. LEOPARD-2, the first multicentre RCT, recruited 99 patients but did not reveal the superiority of the TLPD and ended with an estimated mortality rate of 6%; 5 patients died in the laparoscopic group, and 1 patient died in the open group.¹³ This result limited the application of TLPD and hindered the development of the technique throughout the Netherlands. Therefore, a multicentre-based RCT with available expertise in laparoscopy guaranteeing sufficient enrolment is necessary.

Therefore, in this trial, TJDBPS01, a large registry will be created by collecting data from the fourteen qualified centres in China to create a working basis for analysing outcomes of interest and obtaining evidence for comparing TLPD and OPD. Additionally, it will be possible for TJDBPS01 to acquire long-term outcomes of PD patients who have not been reported in any RCT studies.

Objectives

The broad goal of this trial is to evaluate the safety and efficacy of TLPD and OPD procedures. The main hypothesis is that TLPD has superiority advantages in postoperative recovery and is equivalent in safety to OPD. The primary outcome variable is the postoperative length of stay. The secondary outcome variables include estimated blood loss, operation time, complication rate, R0 resection rate, comprehensive complication index (CCI) and overall survival.¹⁴

METHODS AND ANALYSIS

Design

The TJDBPS01 trial is designed as a multicentre, randomized, controlled, superiority trial with two parallel groups in fourteen pancreatic centres among 11 provinces in China, providing a minimum of >50 PDs annually. **(Supplementary file 1)** All the participating fourteen centres with pancreatic surgery expertise have performed ≥ 104 TLPDs and OPDs, respectively.¹⁵ Screening and identification of eligible patients will take place within the participating centres' pancreatic multidisciplinary team (MDT). All patients about to undergo PD will go through a standard evaluation: Contrast-enhanced multi-thin-slice CT scans (1 mm) with or without endoluminal ultrasound (EUS), which will be discussed by the MDT. Magnetic resonance imaging with diffusion-weighted imaging is applied to rule out the suspected metastasis before surgery. Patients recommended for resection on the basis of a high likelihood of a malignant lesion will be contacted and provided with a detailed consensus. Consent (version date 2018-03-20, Ver 1.0) will be obtained by an investigator who should comply with applicable regulatory requirements and adhere to the ethical principles in the Declaration of Helsinki. It is not necessary for patients to have a histological diagnosis of malignancy before surgery provided that the MDT has confirmed that the lesion is concerning enough to require resection.¹⁶ Randomization will be performed in a 1:1 allocation. The patients will be randomized into two groups: (1) patients who will undergo TLPD and (2) patients who will undergo OPD. This study is structured after the SPIRIT 2013 guideline defining standard protocol items for clinical trials. **(Figure 1)**

Study population

All adult patients with an indication for PD, including but not limited to pancreatic cancer, bile duct cancer, duodenal cancer and benign diseases such as mass-forming pancreatitis, will be evaluated and informed of the possibility to take part in the TJDBPS01 trial. After the consent form is signed, participants will be randomized into the two previously mentioned groups.

Patients and public involvement

Neither patients nor the public are involved in the conception, design or conduct of the study.

Participants eligibility

Inclusion criteria

Inclusion criteria include (1) age between 18 years and 75 years; (2) histologically proven periampullary cancer, including pancreatic cancer, bile duct cancer, and duodenal cancer; (3) cStage I pancreatic cancer; (3) an indication for elective PD surgery; (4) feasibility for both TLPD and OPD (the subject can also undergo surgery according to the MDT); (5) patients understand the nature of this trial and are willing to comply; (6) patients are able to provide written informed consent; and (7) patients are treated with curative intent in accordance with international guidelines.¹⁷

Exclusion criteria

Exclusion factors include (1) distant metastases including peritoneal carcinomatosis, liver metastases, distant lymph node metastases, and involvement of other organs; (2) patients may undergo left, central or total pancreatectomy or palliative surgery other than PD; (3) patients with

high operative risk, as defined by the American Society of Anesthesiologists (ASA), with a score \geq 4; (4) synchronous malignancy in other organs or second cancer requiring resection during the same procedure; (5) pregnancy; or (6) patients underwent or needed neo-adjuvant therapy including medication or radiotherapy.

Interventions

TLPD surgical procedures

The standard operative technique is described as follows (**Figure 2**). Any small variations according to the surgeon's preference, such as a different order for surgical steps, a variation in approach or pancreatic anastomosis technique, will be allowed but must be recorded authentically in the case record form.

All procedures are to be performed by two regularly trained pancreatic surgeons. Patients will be under general anaesthesia and placed in a supine position; an anti-Trendelenburg position (10°-30°) may be used, if necessary. The right arm is placed along the body and the left arm in 90° abduction for accessing arterial monitoring. A 12-mm trocar (VersaportTM, COVIDIENTM) is placed slightly lower than the umbilicus, and pneumoperitoneum is established. Two 12-mm trocars are then placed, both lateral to the first trocar on the right and left in the midclavicular lines. Another two 5-mm trocars are placed at the right and left infracostal arch on both sides of the anterior axillary line (3-4 cm subcostal). Diagnostic laparoscopy is performed to rule out any abnormalities and metastasis. The cystic duct and artery are transected, and the gallbladder is moved and set aside. The round ligament is retracted to the anterior abdominal wall, possibly with either a suture, according to the surgeons' experience. The lesser sac is opened and a Kocher manoeuvre is performed, and the

duodenum is therefore mobilized. The gastro-epiploic artery and vein are transected. The distal stomach is transected on the left side of the pylorus with an endostapler (COVIDIEN™, Endo GIA™, Auto Suture™) after removal of the nasogastric tube. The common hepatic arteries are identified on the inferior border of the pancreas and suspended with a rubber band. Lymph node station 8a is dissected. Subsequently, the right gastric artery and gastroduodenal artery are ligated and transected with at least 2 Hem-o-lok clips (WECK®) and 1 firm ligation (Ethicon®). The portal vein (PV) is then identified at the superior border of the pancreatic neck, and the superior mesenteric vein (SMV) is identified at the inferior border of the pancreas. A tunnel is created posterior to the pancreatic neck and anterior to the SMV and PV. The pancreas is slung with a vascular blocking band and then transected with ultrasonic shears (THUNDERBEAT, OLYMPUS), but the pancreatic duct is transected sharply with scissors and intubated to ensure that it is open. The jejunum is transected approximately 15 cm from the ligament of Treitz with an endostapler (COVIDIEN™, Endo GIA™, Auto Suture™). After retracting the duodenum and jejunum to the right side of the mesenteric root, the duodenum is stretched, and the uncinate process is mobilized. The branches of the SMA and SMV are carefully handled until the uncinate process is fully dissected. Retroportal lymph nodes are resected according to the International Study Group on Pancreatic Surgery guidelines.¹⁸ The common hepatic duct is tunnelled and transected for anastomosis.

Preparation of the pancreatic stump and jejunal loop: The pancreatic stump remnant is dissected to approximately 0.5 cm (no more than 1.0 cm) in length, and careful haemostasis is established with the ultrasonic shears or absorbable sutures. A 6 to 8 Fr plastic catheter is then inserted as a stent into the pancreatic duct remnant to prevent pancreatic duct stenosis after suture placement. The jejunal limb is brought up to the right of the middle colic vessels in a retrocolic manner, and the blind end

is placed near the pancreatic remnant. Negative resection margin status on the frozen section of the specimen will be confirmed before intracorporeal reconstruction if malignancy is suspected.

Pancreaticojejunal anastomosis: An imbedding end-to-side PJ including four layers of mattress sutures is performed.¹⁹ The first layer of the anastomosis is created between the posterior wall of the pancreatic stump and the posterior seromuscular layer of the jejunum. Two completely transpancreatic 4-0 prolene (Premilene, BRAUN) mattress sutures are placed at a point approximately 0.2 cm from the superior and inferior edge of the main pancreatic duct, respectively.

The most critical surgical step in placement of the first and second mattress sutures is interlocking them in the posterior wall of the jejunum. Care should be taken to preset the sutures to avoid passing them through or injuring the main pancreatic duct. Both mattress sutures are preplaced approximately 0.5 cm from the cut edge of the pancreatic remnant. The number of these sutures varies depending on the size of the pancreatic duct but typically ranges between two and four sutures.

After a small, full-thickness jejunotomy is created in line with the pancreatic duct, the second layer of the anastomosis is created between the posterior wall of the pancreatic stump and the posterior wall of the small jejunal hole. Two completely transpancreatic 4-0 prolene mattress sutures are placed following the first two sutures, but a full-thickness mattress suture in the posterior wall of the small hole is made in the jejunum approximately 0.2 cm from the edge. The third and fourth mattress sutures are interlocked in the posterior wall of the jejunal hole. The posterior wall of the jejunal loop is then tightly anastomosed with the posterior wall of the pancreatic stump without any remaining suture intervals. The third layer of the anastomosis is made between the anterior wall of the pancreatic stump and the anterior wall of the jejunum. Two incompletely transpancreatic interlocking 4-0 prolene mattress sutures are placed as described for the second layer of the

anastomosis. However, the mattress sutures do not completely penetrate the pancreatic parenchyma but only enter the tissue half-way. The fifth and sixth mattress sutures are supposed to interlock in the anterior wall of the jejunal hole. The fourth layer of the anastomosis has been made between the anterior wall of the pancreatic stump and the anterior wall of the jejunum. Four incompletely transpancreatic interrupting 4-0 prolene mattress sutures are placed to close the gaps in the superoinferior margin of the anastomosis and strengthen the sutures on its anterior wall, achieving a pancreas-enteric anastomosis. Approximately 10 cm distally, an end-to-side hepaticojjunostomy is performed with running (bile duct dilatation of >5 mm) or interrupted (bile duct dilatation of ≤ 5 mm) sutures (Novosyn, BRAUN). Approximately 40 cm distal to the biliary anastomosis, an antecolic side-to-side gastroenterostomy is performed with the staple technique, and 2 layers of running 3-0 sutures (Novosyn, BRAUN) are used to close the gastric stump. Two 27 Fr surgical drains are placed through Winslow and the upper region in the PJ stoma. After haemostatic control, the trocars are removed. The specimen is extracted through an upper-middle incision, which is subsequently closed in layers together with the closure of the trocars. A video clip for illustrative purpose of the basic laparoscopic surgical procedure is available as **Supplementary file 2** and **Supplementary file 3**.

OPD techniques

The open pancreaticoduodenectomy will be performed by the same group of surgeons. A right sided rectus muscle abdominal incision is preferred for better exposure. The key steps are similar to those taken in the TLPD group. Since outcomes of open pancreatectoduodenectomy worldwide are promising and convincing, the surgical technique in the OPD arm reflects their usual practice; also,

the anastomoses will be performed according to each centre's protocol. Procedure variations according to the surgeon's preferences are allowed but must be recorded on the case record form. Two surgical drains are placed as in the TLPD procedure.

Conversion from TLPD to OPD

Conversion is defined as any TLPD group in which a skin incision is used for reasons other than trocar placement or specimen removal. Subjects allocated to TLPD who undergo intraoperative conversion to open pancreateoduodenectomy will be analysed in the TLPD group according to intention-to-treat principles. Reasons for conversion should be carefully recorded in the surgical record.

Postoperative management

Postoperatively, antibiotics will be administrated as well as either parenteral or enteral nutrition. Somatostatin analogs are not routinely used for preventing postoperative pancreatic fistula. The abdominal drains are kept in place for observation and will be removed if they are not productive and/or the presence of a pancreatic fistula or bile leakage is excluded. Amylase determination of the drains is measured on the third postoperative day or when a fistula is suspected following the ISGPF criteria. Patients are willing to be discharged home when reached criteria as follow: (1) do not need any intravenous infusion, (2) can orally take solid or semisolid food, (3) do not need analgesics or fully comfort by orally analgesics, (4) incision fully recovered without infection, (5) could get off bed without assistance, (6) walking at least 250 meters in plain road, (7) major organ functioning well and nearly normal haematological parameters. The surgeons operating are not involved in the

postoperative management hence can't influence the discharge of the patients.

Follow-up

All patients will be followed-up for a period of 2 years.

Withdrawal

All the patients will be free to participate in this study and can decide to withdraw at any time. If a patient withdraws, his/her information will not be recorded in the study. However, the research team can still collect outcome data from the healthcare records.

Outcomes measures

Primary outcome measure

The primary outcome variable is the postoperative length of stay.

Secondary outcome measures

The secondary outcome variables include the estimated blood loss, operation time, complication rate, mortality, CCI, R0 resection rate, and overall survival.

Missing data

Bias due to missing data will be investigated by comparing the baseline characteristics of participants with and without missing values. Depending on the extent of missing levels, the

predictors of missing values will be identified. The primary outcome analysis will be conducted according to the Intention to treat analysis and will be adjusted for the predictors of missing values as the sensitivity analysis. In addition, multiple imputations will be used to impute missing data, and the imputed data will also be analysed as part of the sensitivity analyses.

Participant timeline

The trial time schedule of enrolment is estimated to be a 2-year period, followed by a 2-year follow-up visit after discharge from the hospital. Once the eligibility of the patients is confirmed, randomization will be applied. When to apply the interventions is not strictly defined: every collaborating centre can make its own preferred adjustment, but this will usually be within 48 hours after the randomization. The assessments and visits for patients are mandatory in the first month, at 3 months, 6 months, 12 months and 24 months, with either telephone or in-hospital follow-up.

(Figure 3)

Sample size

Considering the primary outcome variable is the LOS and according to the latest systemic review,²⁰ MIPD was associated with a shorter length of stay than OPD was (mean difference [MD] -2.95 days, 95% confidence interval [CI] -3.91 to -2.00 days). Accordingly, the sample size is determined based on the primary objective of comparing the LOS between two surgical methods. Based on a mean difference of 2.95 days in the two surgical groups, using the two-sided test with 80% power at a significance level of 5%, the minimal sample size needed to detect a significant difference is calculated to be 274 patients in the TLPD group and 274 in the OPD group ($\alpha=0.05$,

$\beta=0.2$, power=0.8). Considering washout or loss of follow-up, we enlarged the sample size by 20% to compensate for withdrawal/dropout. Then, the final sample size is 656. Thus, there are 328 patients in each group.

The null hypothesis (H0) is that the mean LOS of TLPD group is the same in OPD group. In another word H0: Mean LOSTLPD = Mean LOSOPD. The alternative hypothesis (HA) is that the mean LOS of TLPD group is not the same in OPD group. HA: Mean LOSTLPD \neq Mean LOSOPD. Two sample t-test are used then if the test statistic turns out that $p>0.05$, then it fails to reject the null. The mean LOS of TLPD group is the same in OPD group. Otherwise, if $p<0.05$, the null hypothesis is rejected. The mean LOS of TLPD group is not the same in OPD group. Base on the previous study and the hypothesis, a -2.95 days of mean differences should be obtained.

Recruitment

Each centre will screen eligible patients through the outpatient department or through advertising by the qualified surgeons. The duration of the recruitment period is estimated to be a 24-month interval depending on each centre's recruiting rate. No financial incentives will be provided to trial investigators or patients for enrolment in the recruitment period.

Participating surgeons and centre criteria

To prevent surgeon bias, participating surgeons must satisfy the following criteria: (1) they must have experience in performing more than 104 TLPDs, (2) they must have experience in performing more than 104 OPDs, and (3) they must have completed the Tongji Hospital TLPD training programme.

All the participating centres represent the largest pancreatic surgery centres in their region, ensuring that 50 PDs are performed annually. If willing to join the trial, each surgeon will be mandated to offer two recently unedited surgery videos for evaluation. The video should be sent to the Board and saved appropriately. Only when the Board approves the surgical techniques can the surgeon and his or her centre participate as a collaborator.

Randomization and allocation

We applied the stratified randomized block design for the fourteen centres with a block number of 4, and the design is conceived using SAS statistical software (V. 9.3; SAS Institute, USA). The randomization is centralized through a computer-generated system and performed in a parallel fashion. A data manager who is independent of the data analysis or patient enrolment generated the randomization schedule. Allocation will be announced and handled out by the study coordinators only after a baseline assessment and the patient consenting to participate in the study. The randomization schedule will not be available to study recruiters or physicians. Allocation will be conducted by the independent data manager. Specifically, when appropriate patients are enrolled, the researcher will inform the data manager, and then the random number and exact treatment group will be returned simultaneously.

Blinding

The patients and surgeons are unblinded while the data collectors, outcome assessors, and data analysts are blinded. The primary endpoint of this study is the LOS and is not influenceable by the data collectors, outcome assessors, and data analysts. Since they are blinded and also are not

involved in the pre-, peri- and post-operative management of the patients, thus they have no determination of the LOS. The surgeons only perform the surgery and didn't involve in the postoperative management. The patients have little influence concerning the LOS, since once the discharge criteria are reached, they will be discharge immediately. Most of the criteria are objective conditions like do not need any intravenous infusion, incision fully recovered without infection, major organ functioning well and nearly normal haematological parameters. These conditions won't influenced by the patients even they are unblinded.

Data collection

Patient demographics: year of birth, sex, body mass index, surgical risk (ASA score), comorbidities, previous surgical history, main complaint, preoperative biliary drainage, preoperative blood samples (haemoglobin level, white blood cell count, and granulocyte: lymphocyte (G:L) ratio), plasma level of total bilirubin, carbohydrate antigen (CA) 19-9, CA 125, carcinoembryonic antigen (CEA), alpha-fetoprotein (AFP), date of admission, and income.

Surgery information: operation date, surgical approach (open or total laparoscopic), type of digestive tract reconstruction, anastomosis approach (intracorporeal, extracorporeal), anastomosis performance (linear stapler, circular stapler, hand-sewn or combinations), placement of intra-abdominal drain, placement of nasogastric tube, total operative time, each anastomosis time (pancreaticojejunostomy, cholangiohepaticojejunostomy, pancreatogastrostomy), texture of pancreas, diameter of the main pancreatic duct, estimated blood loss (EBL), intraoperative blood transfusion, conversion to open surgery, intraoperative complications, intraoperative death, and

extent of lymphadenectomy.

Histopathological information: surgical margin status (R0 resection rates), number of retrieved lymph nodes, number of positive lymph nodes, tumour location, size of the tumour, depth of invasion (T classification), lymph node status (N classification), AJCC pathological stage (the eighth edition),²¹ and histological type.²²

Postoperative clinical findings: length of postoperative hospital stays, postoperative blood transfusion, patient mobilization (postoperative day (POD)), liquid diet (POD), soft solid diet (POD), drain removal (POD), length of intravenous analgesic use, cost for hospitalization, cost for surgery, drain production and amylase, haemoglobin levels, white cell count, tumour marker level after surgery, type of complication, reoperation, and Clavien-Dindo grade.²³

Follow-up: date of last follow-up visit, adjuvant chemotherapy, adjuvant radiotherapy, patient status at last follow-up visit (alive, dead, or lost to follow-up), and disease-free or not during follow-up.

All the data will be placed into the local database via a data registry server managed by the data collection group. In this trial, the follow-up and statistical staff will be masked to the intervention arms.

Data management

In this study, the EDC data collection system will be used for data collection. The roles of the staff involved in data collection include the clinical research coordinator (CRC), clinical research associate (CRA), and others.

All staff involved in the data collection process are to be qualified to access the research database. Data collection must be collected in accordance with standard specification processes.

Photographers should work in accordance with the visit process requirements, and completed eCRF will be used to enter the original data into the system with a guide; the system cannot automatically identify the data needed to be entered through manual input by the data entry person.

General statistical methods

SAS statistical software (V. 9.4; SAS Institute, USA), the R program (R Foundation for Statistical Computing, Vienna, Austria, version 3.4.4), Power Analysis and Sample Size (PASS, V11.0 NCSS) and other required statistical software programs will be used, when appropriate. All statistical tests will be tested with two sides, and $P < 0.05$ is considered to represent a statistically significant difference, with a confidence interval of 95%. Quantitative data statistics will be described by the number of cases, means, standard deviation (SD), median, quartile (IQR), maximum and minimum (Min ~ Max) values. Qualitative data statistics will be described by the frequency, constituent ratio, or percentage. The effect research and the analysis comparison will select the data set according to the specific research goal with the appropriate statistical analysis method. The inter-group comparison of qualitative data will use the Chi-square test, and the independent sample T test will be used to compare the quantitative data between the groups.

The inter-group comparison of control covariance can be compared by covariance analysis or the generalized linear model.

We used the SPIRIT checklist when writing this study protocol.^{24,25}

Data monitoring

Efficacy and safety of the duration of data acquisition is 90 days after the patient signs the informed

consent (ICF) and the visit. Security data will evaluate the severity of adverse events based on the CTCAE 4.0 standard. All adverse events will be recorded on the CRF, from the signing of the ICF to the end of the study. LinkDoc acts as the independent third party and will conduct data monitoring throughout the whole trial.

Harms

All the harms, including adverse events or even severe adverse events, will be recorded in detail according to the Common Terminology Criteria for Adverse Events (CTCAE ver 4.0). The data will also be collected by the Hospital's Ethical Committee and in the Chinese Clinical Trial Registry.

Protocol amendments

Any modifications of the protocol that may impact the conduct of the study, potentially benefit the patients or that may affect patient safety, including changes in study design, sample size, and study procedures, will require a formal amendment to the protocol. Additionally, this would need to be submitted to the Hospital Ethics Committee and health authorities must be notified in accordance with local regulations.

Confidentiality

All study-related information and participant information will be stored securely at the study site with locked cabinets in areas with limited access. All local databases will be secured with a password-protected access system.

Publications

Each participating investigator, with equal right, will be able to access the data of the registry, perform statistical analysis, discuss the results and freely write scientific manuscripts. The manuscript must be approved by all the authors before publication.

Ethics and dissemination

Approval from the the Institutional Review Board Approval of Tongji Hospital, Tongji Medical College, Huazhong University of Science and Technology (TJ-IRB20180512) is received in May 2018. All patients signed an informed consent document before entering the study. (**Supplementary file 4**) Consent is obtained by the consultant or designated team member and is preserved by the data collection group. This study has gained ethical approval at both the central and local levels for each participating centers, and central ethical approval has been confirmed by the the Institutional Review Board Approval of Tongji Hospital, Tongji Medical College, Huazhong University of Science and Technology. The recruitment will not start at other centres in the trial until local ethical approval has been obtained. Participating in both groups does not imply any additional risk for the subjects included, since the groups will not be deprived of the application of the most up-to-date recommendations. Any results from this trial (publications, conference presentations) will be published in peer-reviewed journals and conference proceedings.

DISCUSSION

The TJDBPS01 trial is a multicentre, randomized controlled trial. The trial hypothesis is that TLPD

has advantages for postoperative recovery and is equivalent in operation time and oncological results to open pancreaticoduodenectomy. The trial is conducted after the completion of a structured training programme for TLPD in China. The length of the learning curve is analysed, considering the argument on timing. Based on the results of a retrospective study of LPD in China, a minimum of 40 total LPDs is decided as the cut-off point.¹⁵ When the surgeon finished the 104th LPD, the surgical technique would be regarded as mature and stable. Thus, a minimum of 104 TLPDs is decided as the threshold to participate in the TJDBPS01 trial.

As previously mentioned, this is not the only RCT on this topic. Palanivelu et al published their RCT trial results in 2017.¹¹ They conducted a trial comparing LPD and OPD for the treatment of periamppullary tumours in a single tertiary-care teaching institute in India and concluded that laparoscopy offered a shorter hospital stay than open pancreaticoduodenectomy in their randomized trial. The trial recruited 64 patients over a two-year interval period. This study demonstrated a significant advantage for the laparoscopic group in terms of shortened median duration of postoperative hospital stay, which was decreased by half compared with that of open surgery. Additionally, the laparoscopic approach was better regarding blood loss and transfusion requirements in their study, but the difference was less pronounced when procedures with venous resection were excluded. Other outcomes, including the complication rate, lymph node retrieval, and R0 resection, were comparable in the two groups. The authors also reported that their study had limitations, including a relatively small sample size, which resulted in some outcomes not being assessed, such as recurrence and survival (disease-free and overall), and long-term complications. The same conclusion was supported by the Poves study.¹² Unlike the promising conclusions of the previous two studies, the first multicentre trial, LEOPARD-2, demonstrated a concern for the safety

of TLPD.¹³ Their trial found a higher mortality rate in the laparoscopic group, and the authors stated that a small volume centre and an early learning curve might have influenced the outcomes, although all enrolled surgeons were required to have completed training programmes for TLPDs. With all these pros and cons, it is necessary and important to carry out an RCT in a multicentre setting based on the premise that both safety and quality are guaranteed.

In conclusion, the TJDBPS01 trial is a multicentre, randomized controlled trial investigating the safety and effectiveness of TLPD and OPD performed by surgeons who have performed ≥ 104 TLPDs and OPDs, respectively. This trial aims to evaluate TLPD and OPD in daily practice within high-volume pancreatic centres. When this trial is completed with the hypothesis confirmed, it will popularize the application of TLPD and improve patient outcomes.

Acknowledgements We thank Prof. Jingdong Ma and Dr. Tingting Qin from the Department of Epidemiology and Biostatistics, School of Public Health, Tongji Medical College, Huazhong University of Science and Technology, for the statistical support. We thank LinkDoc for the data monitoring. We thank AJE for the English language, grammar, punctuation, spelling, and overall style editing. (**Supplementary file 5**)

Contributors HZ, YCF and JFZ are joint first authors. RYQ, MW and YHY obtained funding for the study. RYQ, HZ and MW designed the study. RFC, XMC, WC, DWL, JDL, HGH, DYL, JL, JHL, JL, YHL, ZJT, WXZ and RYQ performed the operations. XBH and XMY give supervision for the operation. HZ drafted the manuscript. RYQ and MW contributed to critical revision of the manuscript for important intellectual content and approved the final version of the manuscript. All authors have read and approved the final manuscript.

Funding This study is supported by grants from the Hubei Natural Science Foundation (WJ2017Z010), Tongji Hospital Clinical Research Flagship Program 2019CR203.

Competing interests None declared.

Ethics approval is obtained from the Institutional Review Board Approval of Tongji Hospital, Tongji Medical College, Huazhong University of Science and Technology (TJ-IRB20180512). This study has gained ethical approval at both the central and local levels of each participating centers and central ethical approval has been confirmed by the Institutional Review Board Approval of Tongji Hospital, Tongji Medical College, Huazhong University of Science and Technology.

Provenance and peer review Not commissioned; externally peer-reviewed.

REFERENCES

1. Siegel RL, Miller KD, Jemal A. Cancer statistics. CA: a cancer journal for clinicians. 2016; 66(1): 7–30.
2. Tsamalaidze L, Stauffer JA. Pancreaticoduodenectomy: minimizing the learning curve. Journal of visualized surgery 2018;4:64.
3. Gagner M, Pomp A. Laparoscopic pylorus-preserving pancreateoduodenectomy. Surg Endosc 1994, 8: 408–410.

4. Agrusa A, di Buono G, Chianetta D, Sorce V, Citarrella R, Galia M, Vernuccio L, Romano G, Gulotta G. Three-dimensional (3D) versus two-dimensional (2D) laparoscopic adrenalectomy: A case-control study. *Int J Surg.* 2016 Apr;28 Suppl 1:S114-117.
5. Miyamoto R, Inagawa S, Nagai K, Maeda M, Kemmochi A, Yamamoto M. Three-dimension reconstruction of vascular arrangement including the hepatic artery and left gastric vein during gastric surgery. *Springerplus.* 2016 Jun 22;5(1):835.
6. Raspagliesi F, Bogani G, Martinelli F, Signorelli M, Chiappa V, Scaffa C, Sabatucci I, Adorni M, Lorusso D, Ditto A. Incorporating 3D laparoscopy for the management of locally advanced cervical cancer: a comparison with open surgery. *Tumori* 2016; 102(4): 393-397.
7. Chen K, Pan Y, Liu XL, et al. Minimally invasive pancreaticoduodenectomy for perampullary disease: a comprehensive review of literature and meta-analysis of outcomes compared with open surgery. *BMC Gastroenterol* 2017;17:120.
8. de Rooij T, Lu MZ, Steen MW, et al. Minimally Invasive Versus Open Pancreatoduodenectomy: Systematic Review and Meta-analysis of Comparative Cohort and Registry Studies. *Annals of surgery* 2016; 264: 257-267.
9. Nickel F and al. Laparoscopic Versus Open Pancreaticoduodenectomy: A Systematic Review and Meta-analysis of Randomized Controlled Trials. *Ann Surg.* 2019 Apr 8. doi: 10.1097/SLA.0000000000003309. [Epub ahead of print].
10. Zhang H, Wu X, Zhu F, et al. Systematic review and meta-analysis of minimally invasive versus open approach for pancreaticoduodenectomy. *Surgical endoscopy* 2016; 30: 5173-5184.
11. Palanivelu C, Senthilnathan P, Sabnis SC, et al. Randomized clinical trial of laparoscopic

versus open pancreateoduodenectomy for periampullary tumours. *Br J Surg* 2017;104:1443-1450.

12. Poves I, Burdio F, Morato O, et al. Comparison of Perioperative Outcomes Between Laparoscopic and Open Approach for Pancreatoduodenectomy: The PADULAP Randomized Controlled Trial. *Annals of surgery* 2018;268:731-739.

13. van Hilst J, de Rooij T, Bosscha K, et al. Laparoscopic versus open pancreateoduodenectomy for pancreatic or periampullary tumours (LEOPARD-2): a multicentre, patient-blinded, randomised controlled phase 2/3 trial. *Lancet Gastroenterol Hepatol* 2019; 4(3): 199-207.

14. Slankamenac K, Graf R, Barkun J, et al. The Comprehensive Complication Index. A novel continuous scale to measure surgical morbidity. *Ann Surg*. 2013;258:1-7.

15. Wang M, Peng B, Liu J, et al. Practice Patterns and Perioperative Outcomes of Laparoscopic Pancreaticoduodenectomy in China: A Retrospective Multicenter Analysis of 1029 Patients. *Annals of surgery* 2019.

16. Asbun HJ, Conlon K, Fernandez-Cruz L, et al. International Study Group of Pancreatic Surgery. When to perform a pancreateoduodenectomy in the absence of positive histology? A consensus statement by the International Study Group of Pancreatic Surgery. *Surgery*. 2014;155(5):887-892.

17. Nccn clinical practice guidelines in oncology: pancreatic adenocarcinoma. V2. 2019
https://www.nccn.org/professionals/physician_gls/pdf/pancreatic.pdf

18. Tol JA, Gouma DJ, Bassi C, et al. Definition of a standard lymphadenectomy in surgery for pancreatic ductal adenocarcinoma: A consensus statement by the International Study Group on Pancreatic Surgery (ISGPS). *Surgery*. 2014; 156(3):591-600.

19. Wang M, Xu S, Zhang H, Peng S, Zhu F, Qin R. Imbedding pancreaticojejunostomy used in pure laparoscopic pancreaticoduodenectomy for nondilated pancreatic duct. *Surgical endoscopy* 2017;31:1986-1992.
20. Shunda Wang, Ning Shi, Lei You, Menghua Dai, Yupei Zhao. Minimally invasive surgical approach versus open procedure for pancreaticoduodenectomy. A systematic review and meta-analysis. *Medicine* 2018. 96(50): e8619.
21. Chun YS, Pawlik TM, Vauthey JN. 8th Edition of the AJCC Cancer Staging Manual: Pancreas and Hepatobiliary Cancers. *Ann Surg Oncol*. 2018;25(4):845-847.
22. Bosman F, Carneiro F, Hruban R, et al. WHO classification of tumours of the digestive system. 4th edition. IARC, 2010.
23. Clavien PA, Barkun J, de Oliveira ML, et al. The Clavien-Dindo classification of surgical complications: five-year experience. *Ann Surg* 2009;250:187-96.
24. Chan A-W, Tetzlaff JM, Altman DG, Laupacis A, Gøtzsche PC, Krleža-Jerić K, Hróbjartsson A, Mann H, Dickersin K, Berlin J, Doré C, Parulekar W, Summerskill W, Groves T, Schulz K, Sox H, Rockhold FW, Rennie D, Moher D. SPIRIT 2013 Statement: Defining standard protocol items for clinical trials. *Ann Intern Med*. 2013;158(3):200-207.
25. Moher D, Hopewell S, Schulz KF, Montori V, Gotzsche PC, Devereaux PJ, Elbourne D, Egger M, Altman DG: CONSORT 2010 Explanation and Elaboration: Updated guidelines for reporting parallel group randomised trials. *Journal of clinical epidemiology* 2010, 63:e1-37.

Figure Legends

Figure 1 Flow diagram for TJDBPS01.

Figure 2 Representative photographs and laparoscopic views of TLPD. **A:** Kocher manoeuvre; **B:** division of the gastric body; **C:** dissection of the GDA; **D:** division of the pancreatic neck; **E:** dissection of the uncinate process; **F:** picture after specimen removal; **G:** pancreaticojejunostomy; **H:** hepaticojejunostomy; and **I:** gastrojejunostomy.

CBD common bile duct, CHA common hepatic artery, GB gall bladder, GDA gastroduodenal artery,

IVC inferior vena cava, LRV left renal vein, PHA proper hepatic artery, PV portal vein, SMA superior mesenteric artery, SMV superior mesenteric vein.

Figure 3 Standard Protocol Item Recommendations for Interventional Trials (SPIRIT) Figure.

TLPD, total laparoscopic pancreaticoduodenectomy; CT, computed tomography; OPD, open pancreaticoduodenectomy.

Supplementary file 1: The TIF file illustrates the distribution of the collaborators. ■ Heguang Huang from Fujian province, ■ Rufu Chen, Zhijian Tan from Guangdong province, ■ Wei Cheng, Xinmin Yin from Hunan province, ■ Xiaobing Huang, Dewei Li, Jing Li from Chongqing province, ■ Hang Zhang, Yechen Feng, Junfang Zhao, Yahong Yu, Min Wang and Renqi Qin from Hubei province, ■ Jingdong Li from Sichuan province, ■ Deyu Li from Henan province, ■ Jianhua Liu from Hebei province, ■ Jun Liu from Shandong province, ■ Xuemin Chen, Wenxing Zhao from Jiangsu province, ■ Yahui Liu from Jilin province.

Supplementary file 2 and Supplementary file 3: The MP4 files included a video clip of surgical procedures for TLPD.

Supplementary file 4: The PDF file included informed consent in both the Chinese version and the English version.

Supplementary file 5: The PDF file included Editing Certificate for AJE language editing.