

Q-Med AB, a Galderma Affiliate

Clinical Trial No. 43USK1701

STATISTICAL ANALYSIS PLAN

**A Randomized, Controlled, Evaluator-Blinded, Multi-Center Study To Evaluate
The Effectiveness And Safety of *Restylane® Kysse* Versus A Control In The
Augmentation of Soft Tissue Fullness of The Lip**

October 25, 2018

Version 1.0

Revision History:

October 25, 2018 Amendment 1.0

Table of Contents

1	Study Information	4
1.1	Background	4
1.1.1	Study design	4
1.1.2	Number of subjects and randomization	4
1.2	Study objectives	4
1.2.1	Primary efficacy objective	4
1.2.2	Secondary efficacy objectives	4
1.2.3	Safety objectives	5
1.2.4	Exploratory objectives	5
1.3	Efficacy assessments	5
		6
		6
		6
		7
		7
		8
1.4	Efficacy endpoints	9
1.4.1	Primary efficacy endpoint	9
1.4.2	Secondary efficacy endpoints	9
1.4.3	Exploratory efficacy endpoints	10
1.5	Safety assessments	10
1.6	Safety endpoints	10
2	Statistical Methods	12
2.1	General methods	12
2.2	Analysis Populations	12
2.3	Study subjects	12
2.3.1	Subject disposition	12
2.3.2	Protocol deviations	13
2.3.3	Demographic characteristics	13
2.3.4	Medical and surgical history, prior and concomitant medication/procedures	14
2.4	Efficacy analysis	14

2.4.1	Datasets analyzed.....	14
2.4.2	Handling of missing data	14
2.4.3	Primary efficacy analysis	14
2.4.4	Secondary analysis.....	15
2.4.5	Exploratory efficacy analysis.....	16
2.5	Safety Analysis	16
2.5.1	Extent of exposure	16
2.5.2	Adverse events	16
2.5.3	Pre-defined, expected, post-treatment symptoms	17
2.5.4	Safety assessment.....	17
2.5.5	Exploratory safety endpoints	17
2.6	Interim Analysis.....	17
2.7	Determination of Sample Size	18
2.8	Changes in the Analysis Planned in the Protocol	18
3	Reference List.....	19
4	Summary of Changes in SAP Amendment 1.....	20

1 Study Information

1.1 Background

1.1.1 Study design

This is a randomized, controlled, evaluator-blinded, multi-center study to evaluate the effectiveness and safety of *Restylane® Kysse* for lip augmentation and correction of perioral rhytids.

Approximately 280 subjects will be randomized in a 2:1 ratio of treatment to control and treated. Approximately 187 subjects will be treated with *Restylane® Kysse* and 93 subjects will receive the control in the lips and perioral rhytids and will be followed for 48-weeks.

Investigator blinding will be accomplished by using a Treating Investigator to administer the treatments and a Blinded Evaluator, to whom randomization and treatment are concealed, to conduct the blinded assessments.

Safety assessments will be performed by non-blinded personnel. Eligible subjects randomized to receive treatment will be injected by the Treating Investigator at Day 1.

1.1.2 Number of subjects and randomization

Approximately 280 subjects will be randomized to treatment or to control in a 2:1 ratio (treatment: control) stratified by FST (I-III, IV or V-VI) to achieve a total sample size of 234 evaluable subjects (156 randomized to *Restylane® Kysse*, and 78 to *Juvéderm Volbella® XC*). Subjects in the FST=I-III stratum will be further stratified by study centers; subjects in the FST=IV, or FST=V-VI strata will not be further stratified by study centers due to the small sample size.

Before starting the study, a randomization list will be prepared under the supervision of a designated statistician. Randomization will be performed using an Interactive Response System by assigning each subject to *Restylane® Kysse* or the control according to the randomization list.

1.2 Study objectives

1.2.1 Primary efficacy objective

The primary objective of the study is to demonstrate non-inferiority of *Restylane® Kysse* versus a control in lip fullness augmentation by comparing change from baseline in the Blinded Evaluator assessment [REDACTED] at 8 weeks after last injection.

1.2.2 Secondary efficacy objectives

- To evaluate the effectiveness of *Restylane® Kysse* in lip fullness augmentation based on the Blinded Evaluator assessment of MLFS in the upper and lower lip [REDACTED]

- To evaluate the aesthetic improvement of the upper perioral rhytids and oral commissures after treatment with Restylane® Kysse, based on the Blinded Evaluator assessment [REDACTED]
- To evaluate the aesthetic improvement (overall appearance) after treatment *Restylane® Kysse* [REDACTED]
- Subjects' satisfaction after treatment with *Restylane® Kysse* [REDACTED]
- Lip fullness compared to baseline, [REDACTED]

1.2.3 Safety objectives

- To evaluate all AEs at all visits and pre-defined, expected, post-treatment events reported during the first 4 weeks after treatment as recorded in the subject diary.
- To evaluate the safety assessment at all visits, as assessed by a qualified staff member:

- [REDACTED]
- [REDACTED]
- [REDACTED]

1.2.4 Exploratory objectives

- [REDACTED]

1.3 Efficacy assessments

Day 1 is defined as the date of initial treatment.

For all assessments, baseline is defined as the last observation before initial treatment takes place at the baseline visit on Day 1. Change from baseline is defined as the post-baseline value minus the baseline value.

A large black rectangular redaction box covers the majority of the page content, from approximately y=111 to y=284. The box is bounded by a thin white border. There are several smaller black rectangular redaction boxes at the top left (y=111 to y=131), top center (y=111 to y=131), and bottom right (y=261 to y=281).

A large black rectangular redaction box covers the majority of the page content, from approximately y=113 to y=285. The redaction is irregular, with a white rectangular cutout on the right side and a small white square at the bottom right corner.

ation.

Improvement Scale (GAIS)

A 10x10 grid of black bars on a white background. The bars are arranged in a pattern where the width of each bar in a row increases from left to right. The first row has 1 bar, the second has 2, the third has 3, and so on up to the tenth row which has 10 bars. The bars are positioned such that they overlap slightly and are aligned with vertical grid lines.

1.4 Efficacy endpoints

1.4.1 Primary efficacy endpoint

The primary end point is change from baseline in the blinded evaluator assessment

1.4.2 Secondary efficacy endpoints



1.4.3 Exploratory efficacy endpoints

Exploratory efficacy endpoints include:

- (i) Change from baseline in the blinded evaluator assessment

separated by the following subgroups:

- Study site
- Fitzpatrick skin types (I-III, IV and V-VI)
- Race

1.5 Safety assessments

The methods for collecting safety data are described in Section 8 of the Clinical Study Protocol (CSP) and includes the following: assessment of adverse events (AEs) by direct question to subject and evaluation of subject, as well as pre-defined, expected, post-treatment events reported in the subject diaries; evaluation of product palpability, lip texture, symmetry, movement, function and sensation; laboratory assessments (urine pregnancy test only); and device deficiencies.

Each AE, serious as well as non-serious, shall be assessed by the Investigator for causal relationship with the study product and its use (the injection procedure) and for seriousness (Yes or No) of the event.

A two-point scale (Yes or No response) shall be used for the causality assessments. The Investigators shall be asked to indicate a response to each of the following questions in the eCRF:

- “Do you consider that there is a reasonable possibility that the event may have been caused by the study product?”, and
- “Do you consider that there is a reasonable possibility that the event may have been caused by the device injection procedure?”

If any of these questions is answered with a ‘Yes’, the AE will be considered related. Each AE will also be assessed for causal relationship and seriousness by the Sponsor, in order to fulfil regulatory requirements. In case of disagreement, the AE will be classified as “Related”.

1.6 Safety endpoints

Safety endpoints include:

- (i) Incidence, intensity, duration, and onset of related Adverse Events (AEs) collected during the whole study
- (ii) Incidence, intensity, and duration (defined as number of days with the event) of pre-defined, expected post-treatment events collected using a subject diary for 30 days after each treatment and for each treatment area
- (iii) The safety assessment at all visits, as assessed by a qualified staff member:
 - [REDACTED]
 - [REDACTED]
 - [REDACTED]

Exploratory safety endpoints include:

- (i) Incidence and intensity of related AEs, separated by the following subgroups:
 - Study site
 - Injection volume
 - Fitzpatrick skin types (I-III, IV and V-VI)

[REDACTED]

2 Statistical Methods

2.1 General methods

All statistical analyses, including summary tables and data listings, will be performed using the SAS® system (Version 9.3 or higher). Confidence intervals (CIs) will be two-tailed and constructed at a confidence level of 95%. P values will be two-sided and performed at a significance level of 5%.

All study data including observed and derived variables (e.g., change from baseline, response status) used in the summaries of analyses will be presented in by-subject listings. In general, efficacy, safety, demography, subject characteristics, and treatment related variables will be presented using descriptive statistics. For categorical parameters, the number and percentage of subjects in each category will be presented. The denominator for percentage will be based on the number of observations at that time point (i.e. observed cases). For continuous parameters, descriptive statistics will include number of observations, mean, standard deviation (SD), median, and range. Graphs will be used as appropriate.

Any change made to the finalized statistical analysis plan (SAP) after database lock will be documented in the Clinical Study Report (CSR).

For the primary analysis, non-inferiority will only be demonstrated if the CI is entirely below 0.5 for both co-primary endpoints in both the ITT and PP populations. Similarly, superiority will only be demonstrated if the lower confidence limit is entirely below 0 for both co-primary endpoints in both the ITT and PP populations. Therefore no multiplicity adjustment for Type I error is required.

2.2 Analysis Populations

The following populations will be defined for analysis:

- **Safety population:** Includes all subjects who received any of the investigational products, based on the as treated principle (i.e. according to the treatment actually received).
- **Intention-to-treat (ITT) population:** Includes all subjects who were randomized and had a baseline [REDACTED] for both the upper and lower lips. All ITT analyses will be based on the as randomized principle (i.e. according to the treatment they were randomized to).
- **Per protocol (PP) population:** Includes all subjects in ITT who complete the Week 8 visit without any deviations considered to have substantial impact on the primary effectiveness outcome.

The primary efficacy analysis of non-inferiority will be performed both on the ITT and the PP analysis set. Secondary effectiveness and all safety evaluations will be performed based on the ITT and the safety population set, respectively.

2.3 Study subjects

2.3.1 Subject disposition

The number of subjects in each study population (i.e. Safety, ITT and PP) will be summarized. Study population variables will also be presented in a data listing.

The disposition of subjects will be presented by treatment group, and in total, including numbers of subjects that were:

- Screened
- Randomized
- Completed
- Withdrawn (overall and by reason for withdrawal)

All withdrawn subjects will be presented in a by-subject listing.

2.3.2 Protocol deviations

Protocol deviations will be presented in a by-subject listing. Major protocol deviations will be summarized by site and overall.

Depending on the seriousness of the deviation, subject might be excluded from the PP population, which shall be documented prior to database lock.

For this study, the protocol deviations that will exclude subjects from PP are identified (but not limited to) below:

- General
 - Treatment not given according to randomization scheme at initial or touch-up (if touch-up done) treatment
- Efficacy
 - Pre-treatment value missing for at least one lip
 - Any other deviation considered to have substantial impact on the primary effectiveness outcome
- Other
 - Any key inclusion/exclusion criteria not met (with the exception of the ≤ 21 FST V-VI who do not have to meet the Thin/Very thin lip criterion)
 - Use of any prohibited medications or procedures defined in study protocol between initial treatment and the Week 8 visit, that is considered to have a substantial impact on the primary effectiveness outcome

Deviations from the SAP will be documented in the CSR.

2.3.3 Demographic characteristics

Demographic endpoints, baseline assessments, and subject characteristics will be summarized overall and by treatment for the ITT populations.

The following demographic and baseline variables will be included:

- Age (years)
- Sex
- Race
- Ethnicity

- Fitzpatrick skin types (FST)
- Baseline [REDACTED] (assessed by the blinded evaluator and treating investigator)
- Baseline Upper Perioral Lines & Oral Commissures [REDACTED] (assessed by the blinded evaluator and treating investigator)

2.3.4 Medical and surgical history, prior and concomitant medication/procedures

All summaries will be done by treatment group based on the ITT population.

History of surgical events, medical conditions and any prior dermatological procedures or implants will be provided in a by-subject listing.

Prior (with stop dates prior to initial treatment) and concomitant medications (ongoing or with stop dates on or after initial treatment) for all randomized subjects will be provided in a by-subject listing. Prior and concomitant medications will be summarized using World Health Organization Drug Dictionary (WHO-DD) Anatomical-Therapeutic-Chemical (ATC) classification and preferred term (PT) for each treatment. The number and percent of subjects, and the number of drugs, will be summarized by ATC code. Concomitant medications that started due to an AE will be stated in indication.

2.4 Efficacy analysis

2.4.1 Datasets analyzed

The primary efficacy analysis of non-inferiority will be performed both on the ITT and the PP analysis set. Secondary efficacy will be performed based on the ITT population set.

2.4.2 Handling of missing data

Number of missing values will be summarized and reported as appropriate.

Missing data for the primary analysis [REDACTED] will be handled using the hot deck imputation method. Specifically, each missing value [REDACTED] at Week 8 (recipient) will be imputed with a non-missing (observed) value from a subject randomly selected from the same imputation class (donor). Imputation classes will be defined based on treatment group, FST (I-III, or IV-VI) and whether or not the subject met [REDACTED] criterion (Yes, No), [REDACTED]

[REDACTED] missing values will be imputed in a similar manner. Alternative imputation classes may be used if deemed necessary.

Impact of missing data on the primary analysis for Week 8 endpoint will be evaluated by performing sensitivity analysis based on the observed cases in the ITT population.

All other endpoints will be evaluated based on the observed cases in ITT, i.e. no imputations will be done.

2.4.3 Primary efficacy analysis


The primary efficacy analysis is test of non-inferiority of *Restylane® Kysse* to control treatment in the ITT and PP analysis sets, using the confidence interval approach and a non-inferiority margin of 0.5 units. [REDACTED]

Difference between treatments means (control – *Restylane® Kysse*), and the corresponding 2-sided 95% CI, will be calculated and presented for each of the 2 co-primary endpoints using the Student's t-statistic.



2.4.4 Secondary analysis

CIs for responders mentioned below will be calculated using normal approximation.





2.4.5 Exploratory efficacy analysis

- Subgroup analysis of primary efficacy endpoint will be performed following the same method as described for the overall analysis, for the following subsets:
 - Study site
 - Fitzpatrick skin types (I-III, IV and V-VI)
 - Race

2.5 Safety Analysis

2.5.1 Extent of exposure

The number of subjects receiving initial treatment, touch-up treatment, and Week 48 retreatment respectively, will be presented by treatment group and treatment occasion (i.e. initial treatment or retreatment) as well as injection volume and other relevant injection characteristics, such as injection depth, injection method.

2.5.2 Adverse events

AE terms will be coded using MedDRA dictionary. A treatment-emergent AE (TEAE) is defined as an AE with a start date and time on or after the first study product injection. If relationship to treatment is missing, the event will be conservatively summarized as being related to study product. If intensity is missing, a separate category of missing intensity will be included in the summary table, and no imputation of intensity will be performed. Through the data cleaning process, all attempts will be made to avoid missing values for relationship and intensity.

All AEs will be presented in a by-treatment and by-subject listing, detailing the verbatim term given by the investigator, the preferred term (PT), system organ class (SOC), onset date, end date, intensity, outcome, relationship to study product, relationship to study product injection procedure, action taken with study drug, other action taken, affected treatment area, seriousness and criteria for seriousness. Serious AEs will be presented in a separate listing.

Time to onset of an AE will be derived as the start date minus date of most recent treatment. If the start date is missing, it will be assumed that the AE started on the day of most recent treatment.

Duration of an AE will be derived as the stop date minus the start date + 1. If the start date is missing, it will be assumed that the AE started on the day of most recent treatment. Missing stop date will not be imputed and therefore no duration will be calculated in these cases. Instead, the number of AEs that were ongoing at the end of the study will be given.

An overall summary of AEs will be presented by treatment. The summary will include the total number of events, frequency counts and percentages of subjects with:

- Any TEAE

- Any AE related to product and/or injection procedure
- Unrelated TEAEs
- No TEAEs reported

Summaries of related AEs (including the total number of events, frequency counts and percentages of subjects) will be displayed by treatment and treatment occasion according to the following:

- All AEs by SOC and PT in descending order of frequency (the combined frequency of both treatments)
- All AEs by SOC, PT, and maximum intensity (mild, moderate, or severe)
- All AEs by SOC, PT, and action taken (none, medical treatment, non-pharmacological treatment, subject withdrawn)

For related AEs, the number of days to onset and duration of event will be summarized by SOC and PT, using mean, SD, minimum, maximum, and median statistics.

2.5.3 Pre-defined, expected, post-treatment symptoms

Frequency counts and percentages of subjects reporting each pre-defined, expected, post-treatment symptoms (i.e. bruising, redness, tenderness, swelling, pain, itching, lumps/bumps, discoloration, and other symptoms), as collected in the 30 day diary after each treatment, will be presented in total and by maximum severity, for each treatment area separately.

Number of days with the event will also be summarized using mean, SD, min, max, and median.

2.5.4 Safety assessment

will be presented with frequency counts and percentages of subjects classified as “Normal” or “Abnormal” by visit.

Urine pregnancy test results will be presented in a by-subject listing.

2.5.5 Exploratory safety endpoints

- Related AEs by SOC, PT and maximum intensity will be separately summarized for the following subgroups:
 - Study site
 - Injection volume
 - Fitzpatrick skin types (I-III, IV and V-VI)
- [REDACTED]

2.6 Interim Analysis

There is no interim analysis planned.

2.7 Determination of Sample Size

Accounting for approximately 15% drop-outs and non-evaluable subjects due to protocol deviations at week 8, a total of approximately 280 subjects will be randomized in the study. It is expected that the number of randomized subjects will be similar across the study sites.

2.8 Changes

3 Reference List

4

