

# Safety of TofAcitinib in Routine Care Patients With Rheumatoid Arthritis (STAR-RA) - Cardiovascular Endpoints

May 19, 2021

NCT04772248

## 1. Comparison Details

### a. Intended aim(s)

- To compare the risk of cardiovascular events (including myocardial infarction or stroke), between patients treated with tofacitinib and patients treated with TNF inhibitors (TNFi) for rheumatoid arthritis (RA) among, 1) “real world evidence (RWE)” cohorts including routine care patient population from the US and, 2) “Randomized controlled trial (RCT) DUPLICATE” cohorts including routine care patient population who meet inclusion and exclusion criteria of the *Safety Study Of Tofacitinib Versus Tumor Necrosis Factor (TNF) Inhibitor In Subjects With Rheumatoid Arthritis* (“ORAL Surveillance”, NCT02092467) clinical trial.
- To examine the risk of myocardial infarction, stroke, heart failure hospitalization, coronary revascularization, and all-cause mortality as separated endpoints when comparing tofacitinib with TNFi in patients with RA among, 1) “real world evidence (RWE)” cohort including routine care patient population from the US and, 2) “Randomized controlled trial (RCT) DUPLICATE” cohort including routine care patient population who meet inclusion and exclusion criteria of the *Safety Study Of Tofacitinib Versus Tumor Necrosis Factor (TNF) Inhibitor In Subjects With Rheumatoid Arthritis* (“ORAL Surveillance”, NCT02092467) clinical trial.

### b. Primary endpoint

- Major adverse cardiovascular events (MI and stroke)

## 2. Person responsible for implementation of analysis

Hemin Lee, MD, MPH will implement the study design and analysis in the Aetion Evidence Platform. She is not responsible for the validity of the design and analytic choices. All implementation steps will be recorded and the implementation history will be archived in the platform. Propensity score (PS) fine stratification weighting will be implemented using SAS statistical software (SAS Institute, Cary, NC).

## 3. Data Source(s)

US MarketScan, 2012-2018

Optum, 2012-2020

Medicare Claims Database, 2012-2017

#### 4. Study Design Diagrams

**Figure 1. Inclusion and Exclusion Criteria for RWE and RCT Duplicate Cohorts**

##### **Real World Evidence Cohorts**

###### **Inclusion Criteria**

- Patients treated with tofacitinib or TNF inhibitors in MarketScan (2012-2018), Optum (2012-2020), and Medicare fee-for-service (Parts A, B and D; 2012-2017)
- At least 365 days of continuous enrollment in health plan prior to and including cohort entry date
- At least 2 RA codes between 7 and 365 days apart

###### **Exclusion Criteria**

- Index drug in 365 days prior to cohort entry date (prevalent users)
- < 18 years of age (Optum, MarketScan), < 65 (Medicare)
- Missing data on age or gender
- Nursing home or hospice admission
- TNFi users with prior prescriptions of JAK inhibitors (tofacitinib, upadacitinib, or baricitinib)
- TNFi users initiating with multiple TNFi on same date
- Tofacitinib users with prior prescriptions of baricitinib, upadacitinib
- Tofacitinib users initiating treatment with tofacitinib and baricitinib (or upadacitinib) on same day

##### **RCT-Duplicate Cohorts**


###### **Additional Inclusion Criteria**

- Patients with at least one prescription of methotrexate
- Patients with at least one cardiovascular risk factor (including smoking, hypertension, dyslipidemia, diabetes mellitus, ischemic heart disease, family history of ischemic heart disease)


###### **Additional Exclusion Criteria**

- < 50 years of age (Optum, MarketScan), < 65 (Medicare)
- Patients recently hospitalized with infections
- Pregnant patients

**Figure 2: Study Design for RWE Cohorts**



**Figure 3: Study Design for RCT-Duplicate Cohorts**



## 5. Cohort Identification

### a. Cohort Summary

This study will employ an active comparator, new user observational cohort study design comparing initiators of tofacitinib to tumor necrosis factor inhibitors (infliximab, adalimumab, certolizumab pegol, etanercept, and golimumab).<sup>1</sup> The patients will be required to have continuous enrollment in their health plan during the baseline period of 365 days before initiation of tofacitinib or TNFi (cohort entry date). Follow-up begins the day after drug initiation.

There will be two independent study populations:

- 1) Real-World Evidence (RWE) cohorts: This study population will reflect the patients diagnosed with RA who are routinely treated and managed in setting of clinical practice.
- 2) RCT-duplicate cohorts: This study population will emulate “the Safety Study Of Tofacitinib Versus Tumor Necrosis Factor (TNF) Inhibitor In Subjects With Rheumatoid Arthritis” clinical trial (“ORAL Surveillance”, **NCT02092467**). The inclusion and exclusion criteria of this RCT will be applied to this study population.

### b. Inclusion and Exclusion Criteria Common to RWE and RCT-duplicate Cohorts

*Cohort entry date:*

- First TNFi or tofacitinib dispensation/administration date

*Inclusion criteria:*

- Patients treated with tofacitinib or TNF inhibitors in IBM MarketScan (2012-2018), Optum (2012-2020), and Medicare fee-for-service (Parts A, B and D; 2012-2017)
- A minimum of 365 days of continuous enrollment in health plan prior to (and including) the cohort entry date
- Two diagnosis codes for RA in any setting in 365 days baseline period (diagnosis codes between 7 and 365 days apart)

*Exclusion criteria:*

- Index drug in 365 days prior to cohort entry date (prevalent users)
- Missing data on age or gender
- Admission to nursing facility or hospice on or prior to cohort entry date (ever look-back)
- TNFi users with history of any Janus kinase (JAK) inhibitors (tofacitinib, upadacitinib, or baricitinib) (ever look-back period)
- TNFi users initiating with more than one TNFi on same date
- Tofacitinib users with a prescription of baricitinib, upadacitinib (ever look-back period)
- Tofacitinib users initiating treatment on multiple JAK inhibitors on same day (tofacitinib and baricitinib, tofacitinib and upadacitinib)

**c. Exclusion criteria specific to RWE cohorts**

- Patients less than 18 years of age (MarketScan and Optum) and 65 years of age (Medicare) at cohort entry

**d. Inclusion criteria specific to RCT-duplicate cohorts**

- Patients with at least one methotrexate dispensation (six months look-back period)
- Patients with at least one cardiovascular risk factor (including smoking, hypertension, dyslipidemia, diabetes mellitus, ischemic heart disease, family history of ischemic heart disease) (one-year look-back period)

**e. Exclusion criteria specific to RCT-duplicate cohorts**

- Patients less than 50 years of age (MarketScan and Optum) and 65 years of age (Medicare) at cohort entry
- Patients recently hospitalized with infections (30-day look-back period)
- Pregnant patients (one year look-back period)

## 6. Variables

### a. Exposure-related variables:

#### Study drug:

The study exposure of interest is initiation of tofacitinib

#### Comparator:

Initiators of tofacitinib will be compared to initiators of TNFi (infliximab, adalimumab, certolizumab pegol, etanercept, and golimumab)

### b. Covariates (assessed during 365 day baseline period, unless otherwise specified):

Demographics and Lifestyle Factors	
Age on the cohort entry date	Calendar year of cohort entry
Gender	Obesity
Race	Smoking
Rheumatoid arthritis related factors	
csDMARDs use (methotrexate, hydroxychloroquine, leflunomide, sulfasalazine)	Number of distinct csDMARDs
Glucocorticoids use (365 day baseline, within 60 days of baseline, cumulative dose)	Number of distinct bDMARDs
CVD factors	
Atrial Fibrillation	Coronary Artery Disease
Type 2 Diabetes	Heart Failure
Stroke or transient ischemic attack	Hypertension
Hyperlipidemia	Peripheral vascular disease
Venous thromboembolism	



Comorbid conditions	
Cancer (except non-melanoma skin cancer)	Inflammatory bowel disease
Chronic liver disease	Psoriasis
Chronic kidney disease (stage 3+)	Combined comorbidity index
Chronic obstructive pulmonary disease	Frailty score
Comedication use	
Antidepressant drugs (including serotonin-norepinephrine reuptake inhibitors, selective serotonin reuptake inhibitors, tricyclic antidepressants)	Antihypertensive drugs (including angiotensin II receptor blockers, angiotensin converting enzyme inhibitors, beta blockers, calcium channel blockers, diuretics, nitrates)
Non-steroidal anti-inflammatory drugs	Lipid lowering drugs (Statins and non-statins)
Selective Cox-2 inhibitors	Anti-arrhythmic drugs
Opioids	Anticoagulants
COPD drugs (bronchodilators, inhaled corticosteroids)	Antiplatelets (including acetylsalicylic acid)
Other hormonal agents/HRT	Non-insulin diabetes medications
	Insulin
Markers for healthy behavior, frailty, healthcare use	
Number of C-reactive protein tests ordered	Echocardiogram
Number of serum creatinine tests ordered	Cardiac stress test
Number of lipid tests	Colonoscopy
Number of outpatient visits	Flu Shot
Number of primary care physician visits	Mammography
Number of rheumatologist visits	Pap smear
Number of emergency department visits	Pneumococcal vaccine
Number of hospitalizations (any, recent)	PSA test
Number of hospital days	Total distinct brand drugs used
Cardiologist visits	Total distinct generic drugs used
Electrocardiogram	Number of unique prescription medications

## 7. Outcome variables and study follow-up:

- **Primary outcome:** Composite cardiovascular adverse events (myocardial infarction, stroke). Outcome will be defined by one inpatient claim.
- Secondary outcomes: Inpatients claims for myocardial infarction, stroke, heart failure hospitalization, coronary revascularization (including coronary artery bypass grafting and percutaneous coronary intervention), all-cause mortality

Outcome	Codes	Care Setting	Position
Myocardial infarction	<u>ICD-9 DX:</u> 410.** (except 410.*2)  <u>ICD-10 DX:</u> I21.0-I21.4, I22.*	Inpatient	Primary/ Secondary
Stroke	<u>ICD-9 DX:</u> 430.**, 431.**, 433.*1, 434.** (excluding 434.x0), 436.**  <u>ICD-10 DX:</u> I60.**, I61.*, I63.***	Inpatient	Primary
Heart failure hospitalization	<u>ICD-9 DX:</u> 398.91, 402.*1, 404.*1, 404.*3, 428.**  <u>ICD-10 DX:</u> I09.81, I11.0, I13.0, I13.2, I50.**	Inpatient	Primary
Coronary revascularization	<u>ICD-9 PX:</u> 00.66, 36.03-36.34, (except 36.04)  <u>ICD-10 PX:</u> 0210*, 0211*, 0212*, 0213*, 021K0Z5, 021K4Z5, 021L0Z5, 021L4Z5, 0270***, 0271***, 0272***, 0273***, 02C0*Z6, 02C2*Z6, 02C00ZZ, 02C03ZZ, 02C04ZZ, 02C10ZZ, 02C1*Z6, 02C13ZZ, 02C14ZZ, 02C20ZZ, 02C23ZZ, 02C24ZZ, 02C3*Z6, 02C30ZZ, 02C33ZZ, 02C34ZZ, 02QA0ZZ, 02QB0ZZ, 02QC0ZZ, 02QA3ZZ, 02QA4ZZ, 02QB3ZZ, 02QB4ZZ, 02QC3ZZ, 02QC4ZZ  <u>CPT-4:</u> 33140, 33141, 33510-33536, 33545, 33572, 92920, 92921, 92924, 92925, 92928, 92929, 92933, 92934, 92937, 92938, 92941, 92943, 92944, 92973, 92980, 92981, 92982, 92984, 92995, 92996	Inpatient	Any

**Safety of TofAcitinib in Routine care patients with Rheumatoid Arthritis (STAR-RA)-Cardiovascular Endpoints Protocol**

All-cause mortality		Inpatient (Optum and MarketScan) Any setting (Medicare)	
---------------------	--	--	--

## 8. Study Follow-up Period:

For the primary analysis, we will implement an as-treated exposure definition. Follow-up will start the day after initiation of tofacitinib and TNFi and will continue until the earliest date of the following events:

- The first occurrence of the outcome of interest
- The date of end of enrollment in the database
- End of the study period
- Date of treatment switch or drug discontinuation, defined as the date of the last continuous treatment episode of the index drug plus a defined grace period (i.e., discontinuation defined after 60 days of no prescription refills for the index exposure after the days supply end for the most recent dispensing)
- Death

For secondary analysis, we will implement, an *intention-to-treat* exposure definition. Follow-up will start the day after initiation of tofacitinib and TNFi and will continue until the earliest date of the following events:

- The first occurrence of the outcome of interest
- The date of end of enrollment in the database
- End of the study period
- Death
- 365 days after the cohort entry date

## 9. Propensity score fine stratification analysis

We will use a propensity-score (PS)-based approach to account for measured confounding in this study.<sup>2</sup> The PS will be calculated as the predicted probability of initiating the exposure of interest (i.e., tofacitinib) conditional on baseline covariates using multivariable logistic regression constructed separately in each data source.<sup>3</sup> On average, patients with similar PSs have similar distribution of potential confounders used to estimate the PS. We will trim non-overlapping regions of the propensity score distributions.<sup>3</sup> Subsequently, we will use PS fine stratification weighting to achieve covariate balance between treatment groups.<sup>4</sup> We will create 50 PS strata, based on the distribution of PS in tofacitinib-treated patients. A weight of one will be assigned to tofacitinib users and TNF inhibitor initiators will be weighted proportional to the distribution of tofacitinib initiators in the PS stratum into which they fell to achieve covariate balance ( $N_{\text{exposed in PS stratum } i} / N_{\text{total}} / (N_{\text{reference in PS stratum } i} / N_{\text{total reference}})$ ).<sup>4</sup> This weighting strategy estimates the average treatment effect on the treated (ATT).<sup>4</sup>

We will report multiple diagnostics for PS analysis in this protocol. First, the PS distribution overlap will be provided between two groups before weighting.<sup>4</sup> The balance in each individual covariate between two treatment groups will also be reported before and after weighting using standardized differences.<sup>5,6</sup> Distribution of PS weights will be assessed and truncation strategies will be considered if necessary.

## 10. Statistical analysis plans

Descriptive statistics, including frequencies and proportions for categorical variables and mean, median, standard deviation (SD), interquartile range (IQR), and range (minimum, maximum) for continuous variables, will be used to summarize baseline characteristics for each study cohort. Standardized differences will be used to compare characteristics before and after PS fine stratification weighting.<sup>5,6</sup> Crude incidence rates and corresponding 95% confidence intervals (CI) will be reported for each study outcome. The primary analysis will consist of Cox proportional hazards model weighted using PS fine stratification weights and using an *as-treated* exposure definition. This model will generate weighted hazard ratio and 95% confidence interval when comparing tofacitinib with TNFi. We will also report difference in rates when comparing tofacitinib with TNFi and corresponding 95% CI in the weighted population. Results will be reported independently for each database. Effect estimates will also be pooled across three databases using fixed effects model.

We will also report cumulative incidence and the corresponding 95% CI independently for all outcomes and each study group. Pre-specified subgroup analysis will be conducted based on age ( $\leq 65$  and  $> 65$ ), sex, and baseline cardiovascular disease (RWE cohorts only). Secondary analysis will be conducted by using an *intention-to-treat* exposure definition whereby patients will be censored 365 days after initiation of treatment with tofacitinib or TNF inhibitors. We will also conduct propensity score matching where each patient initiating tofacitinib will be matched with those initiating TNFi using nearest

neighbor greedy matching without replacement using a caliper of 0.025 on the natural scale of the PS (1:1 matching).<sup>3,7</sup> Standardized differences will be used to compare characteristics before and after matching. In addition, we will examine the risk of cardiovascular outcomes and all-cause mortality by stratifying by number of previous prescriptions of biological DMARDs (no previous prescriptions of bDMARDs vs  $\geq 1$  prescription of bDMARDs). Finally, sensitivity analyses will be conducted by restricting the TNFi comparator group to only adalimumab and etanercept users and by using herpes zoster as a positive control outcome.<sup>8</sup>

## 11. References

1. Lund JL, Richardson DB, Sturmer T. The active comparator, new user study design in pharmacoepidemiology: historical foundations and contemporary application. *Curr Epidemiol Rep*. 2015;2(4):221-228.
2. Rosenbaum PR RD. The central role of the propensity score in observational studies for causal effects. . *Biometrika*. 1983;70(1):41-55.
3. Austin PC. An Introduction to Propensity Score Methods for Reducing the Effects of Confounding in Observational Studies. *Multivariate Behav Res*. 2011;46(3):399-424.
4. Desai RJ, Franklin JM. Alternative approaches for confounding adjustment in observational studies using weighting based on the propensity score: a primer for practitioners. *BMJ*. 2019;367:l5657.
5. Franklin JM, Rassen JA, Ackermann D, Bartels DB, Schneeweiss S. Metrics for covariate balance in cohort studies of causal effects. *Stat Med*. 2014;33(10):1685-1699.
6. Austin PC. Using the Standardized Difference to Compare the Prevalence of a Binary Variable Between Two Groups in Observational Research. *Communications in Statistics - Simulation and Computation*. 2008;38(6):1228-1234.
7. Austin PC. A comparison of 12 algorithms for matching on the propensity score. *Stat Med*. 2014;33(6):1057-1069.
8. Pawar A, R D, N G, et al. Risk of admission to hospital for serious infection after initiating tofacitinib versus biologic DMARDs in patients with rheumatoid arthritis: a multidatabase cohort study. *Lancet Rheumat*. 2020;2(2):e84-e98.