



**SERVIZIO SANITARIO REGIONALE
EMILIA - ROMAGNA**
Istituto Ortopedico Rizzoli di Bologna
Istituto di Ricovero e Cura a Carattere Scientifico



“Global collaboRAtion on CIC-DUX4, BCOR-CCN3, high-grade Undifferentiated round cell sarcoma (URCS) project”

Study code	GRACefUI
Sponsor's Name and Address:	IRCCS Istituto Ortopedico Rizzoli Via di Barbiano 1/10 40136 Bologna Italy
Study Number/Version/Date:	Vers 3.0 06 Feb 2023
Coordinating Center:	IRCCS Istituto Ortopedico Rizzoli SC Osteoncologia sarcomi dell'osso e dei tessuti molli e terapie innovative Via Pupilli 1 40136 Bologna, Italy
Coordinating Investigator and address:	Emanuela Palmerini MD Phone: 051-6366.199 Email: emanuela.palmerini@ior.it
Methodology:	Experimental study with biological material : case series review of clinical and histological data
Type:	Academic
Founding:	None

Protocol Signature

Sponsor: IRCCS Istituto Ortopedico Rizzoli 40136 Bologna, Italy

Study Title: “Global collaborAtion on CIC-DUX4, BCOR-CCN3, high-grade Undifferentiated round cell sarcoma (URCS) project’

Study Acronym/Code: GRaCefUI

Protocol Version and Date: Version 3.0 of 06 Feb 2023

I read this protocol and I accept to conduct this trial in accordance with the protocol stipulations, GCP guidelines and the Declaration of Helsinki.

Study global coordinator :

Dott. ssa Emanuela Palmerini
SC Osteoncologia sarcomi dell’osso e dei tessuti molli
e terapie innovative
IRCCS Istituto Ortopedico Rizzoli
40136 Bologna, Italy

Local Site Name: IRCCS Istituto Ortopedico Rizzoli

Local Site Principal Investigator Name: Dr. Emanuela Palmerini

Local Site Principal Investigator Signature

Signature Date

Emanuela Palmerini

06 Feb 2023



TABLE of CONTENTS

Background	pag. 5
Study Design	pag. 5
Objective of the study	pag. 5
Population	pag. 6
Material and Methods	pag. 6
Statistics	pag. 7
Enrolment procedure	pag. 7
Data Collection	pag. 8
Ethic and quality assurance	pag. 8
Informed Consent	pag. 8
Confidentiality	pag. 9
Publication of the results	pag. 9
Sponsor Role and Responsibility	pag. 10
References	pag. 11

BACKGROUND

Undifferentiated round cell sarcomas (URCSs) represent a diagnostic challenge. URCSs cell histology and expression of CD99 mostly resemble Ewing sarcoma. However, they also may include other tumors such as mesenchymal chondrosarcoma, synovial sarcoma, or small cell osteosarcoma. Definitive classification usually requires detection of entity-specific mutations.

Recently, the use of modern molecular techniques led to the description of a variety of new, rare, sarcoma subtypes, which have morphological appearance similar to Ewing sarcomas ("Ewing-like" sarcomas), but carry gene fusions different from that of Ewing sarcoma, with CIC-DUX4, and BCOR-CCNB3 being the most frequent. Other non-Ewing round cells sarcomas are characterized by rearrangement of CIC (CIC-rearranged sarcomas) or BCOR (BCOR-rearranged sarcomas) with other genes, by Internal Tandem Duplication of BCOR or by other translocations (i.e.: CRTC1-SS18).

The identification of these gene fusions suggests that other, yet to be identified, gene fusions could be associated with this type of tumour. Furthermore, novel gene fusions in Ewing-like" sarcomas have been reported recently, in case reports, including CIC-FOXO4 BCOR-MAML3 and ZC3H7B-BCOR. Identifying these genetically defined entities may contribute towards understanding the pathogenesis and the behaviour of these tumours. The optimal treatment for these ultra rare tumors is not known.

We propose to collect all cases of URCSs treated within the referral centres for sarcoma, including Italian sarcoma group network (ISG), in order to reclassify and describe their treatment and outcome.

STUDY DESIGN

This is a multicenter retrospective and prospective study that will analyze all cases with a diagnosis of URCSs from 01 January 1983 and all new cases referring to each participant site.

OBJECTIVE OF THE STUDY

Primary Objective

Correlate clinical and pathological variables with event-free survival (EFS) and overall-survival (OS) in URCSs

Secondary objective

Molecular re-classification of URCSs based on presence of CIC-rearrangement, and BCOR-CCNB3 fusion genes or of other genetic alterations.

Exploratory objective

1. In all the “true-negative” URCSs (unclassified URCS), genomic/genetic characteristics will be correlated with clinical outcome.
2. In all URCS samples with frozen tissue available, potential diagnostic and therapeutic target will be searched through RNA-seq

POPULATION

Inclusion criteria

1. Patients with URCSs treated from 01 January 1983 and all new patients will be included.
The study will include the following:
 - CIC –rearranged sarcomas
 - BCOR-CCNB3 fusion-positive sarcomas and sarcomas with BCOR genetic alterations
 - Undifferentiated round cell sarcoma with other genetic alterations
 - Undifferentiated round cell sarcomas with no evidence of genetic alterations (unclassified URCS)
2. Clinical and outcome information available.
3. Histological slides/formalin-fixed paraffin-embedded tissue tumor (FFPE) blocks from archive available to perform the histology analysis/frozen tissue representative of the tumor available.

Exclusion criteria

- 1) Diagnosis different from URCSs
- 2) Patient with no clinical or outcome information available

The study will include male and female patients without limit of age

MATERIAL AND METHODS

Patients with a morphological diagnosis of URCSs will be included and will be classified as

- CIC –rearranged sarcomas
- BCOR-CCNB3 fusion-positive sarcomas and sarcomas with BCOR genetic alterations
- Undifferentiated round cell sarcoma with other genetic alterations
- Undifferentiated round cell sarcomas with no evidence of genetic alterations (unclassified URCS)

Formalin-fixed paraffin embedded tissue and/or frozen tumoral tissue of URCSs cases *negative* for all Ewing’s sarcoma molecular fusions, will be analyzed for the presence of CIC and BCOR-CCNB3 translocations fusion at each Institution. All these analyses will be performed by using real-time PCR, FISH analysis or NGS.

Positive cases will be then re-classified accordingly, while negative cases will be defined as “true-negative”. For patients defined as “true-negative, NGS or RNA-seq will be performed (if not already done), in order to detect the possible presence of other genetic alterations.

Cases that remain negative for genetic alterations will be labelled as unclassified URCS.

Treatment, outcome, and prognostic factors will be related with the molecular re-classification.

Each participating center will have to send to the global study coordinator the *pathology form* correctly filled in.

RNA-seq is an increasing used methodology to study either coding and non-coding RNA expression, for the study of the transcriptome based on next-generation sequencing (NGS). In all URCS samples with frozen tissue available, differentially expressed genes, and potential diagnostic and therapeutic targets will be searched through RNA sequencing

STATISTICS

For survival analysis the following factors will be correlated with Overall Survival (OS) and Event Free Survival (EFS): stage (localised vs metastatic), histology (CIC-DUX4, BCOR-CCNB3, other), age (pediatric < 18 years vs adult ≥ 18 years), gender, pathologic response (good vs poor) to neoadjuvant chemotherapy.

EFS and OS will be estimated according to the Kaplan and Meier method with their respective 95% confidence intervals (CI) and calculated from the first day of chemotherapy administration to

death or last follow-up visit. A Cox regression model will also be used to examine the effect of factors on OS and EFS.

For retrospective analysis, it is expected to include about 200 patients. For prospective study, we expect to include 60 patients.

ENROLLMENT PROCEDURE

Patients considered eligible and who have provided a written informed consent will be included into the study and their material will be used for the purpose of this research according to the protocol and to the informed consent provided.

For the retrospective analysis, the investigator will extract from Institute database the list of potential eligible patients for whom the tumor material is available. After the verification of the presence of the study enrollment criteria, the patient will be included in the study.

Due to the high incidence of mortality of the disease under investigation, it would be possible that some eligible subjects will be deceased.

DATA COLLECTION

Clinical data will be retrieved by reviewing clinical charts, radiology and pathology report by local investigators.

A protocol-specific CRF reporting pathologic report at the time of study inclusion and the results of re-classification will be provided.

A CRF is required and should be completed for each included subject.

ETHICS AND QUALITY ASSURANCE

The clinical trial protocol and its documents will be sent before initiating the study to the competent Authorities and Ethics Committees of each participating country for its approval.

The responsible investigator will ensure that this study is conducted in agreement with either the most updated Declaration of Helsinki and all the international and local laws that apply to clinical trials and to patient protection.

The protocol has been written, and the study will be conducted according to the principles of the ICH Harmonized Tripartite Guideline for Good Clinical Practice

(ref: <http://www.emea.eu.int/pdfs/human/ich/013595en.pdf>).

INFORMED CONSENT

All patients will be informed, by the investigator, of the aims of the study, the possible risks and benefits that will derive from the study participation.

The Investigator must clearly inform that the patient is free to refuse participation in the study and that can withdraw consent at any time and for any reason.

They will be informed as to the strict confidentiality of their patient data, but that their medical records may be reviewed for trial purposes by authorized individuals other than their treating physician.

The informed consent procedure must conform to the ICH guidelines on Good Clinical Practice. This implies that "the written informed consent form should be signed and personally dated by the patient or by the patient's legally acceptable representative".

The Investigator must also sign the Informed Consent form, and will keep the original at the site and a copy of the original must be handed to the patient.

The competent ethics committee for each Institution participating to the study must validate local informed consent documents before the study can be opened. It will be emphasized that the participation is voluntary and that the patient is allowed to refuse further participation in the study whenever he/she wants. This will not prejudice the patient's subsequent care.

For retrospective analysis, due to the high incidence of mortality of the disease under investigation, it would be possible that some potential eligible subjects will be deceased.

CONFIDENTIALITY

In order to ensure confidentiality of clinical trial data as disposed the national and European applicable regulation, data will be only accessible for the trial Sponsor and its designees, for monitoring/auditing procedures, the Investigator and collaborators, the Ethics Committee of each corresponding site and the Health Authority.

Investigator and the Institution will allow access to data and source documentation for monitoring, auditing, Ethic Committee revision and inspections of Health Authority, but will have to maintain at all times personal data confidentiality as specified in all the applicable regulations.

All patients included in the study will be identified with a numeric code. Study Responsible and Collaborators of each laboratory and participating Institutions will have to keep a patients' study

inclusion registry containing the following personal data of each patient: name, surname, address and corresponding identification code used for this study.

Study Responsible, Investigators and Collaborators of each laboratory and participating Institutions will guarantee that patients anonymity will be kept at all times and that their identity will be protected from access from unauthorized persons and Institutions.

PUBBLICATION OF RESULTS

The results from this study can be published or shown at scientific conferences. According to usual practice, this multicenter study will be published as a whole, and not with the data obtained separately from each of the sites participants. It is expected that other articles are published about the exploratory aspects of this trial once the main data has been published. The final publication of the trial results will be written by the Coordinator Investigator.

All publications (papers, abstracts, presentations...) including data from the present trial will be submitted for review to all co-authors prior to submission.

SPONSOR ROLE AND RESPONSIBILITY

The sponsor is the sole owner of the data and is responsible of all the clinical trial activities from study design, development, data collection, management, analysis, interpretation of data, writing and the decision to submit the report for publication written by the Principal Investigator.

REFERENCES

1. Gambarotti M, Benini S, Gamberi G, Cocchi S, Palmerini E, Sbaraglia M, Donati D, Picci P, Vanel D, Ferrari S, Righi A, Dei Tos AP. CIC-DUX4 fusion-positive round-cell sarcomas of soft tissue and bone: a single-institution morphological and molecular analysis of seven cases. *Histopathology*. 2016 Oct;69(4):624-34.
2. Italiano A, Sung YS, Zhang L, Singer S, Maki RG, Coindre JM, Antonescu CR. High prevalence of CIC fusion with double-homeobox (DUX4) transcription factors in EWSR1-negative undifferentiated small blue round cell sarcomas. *Genes Chromosomes Cancer*. 2012 Mar;51(3):207-18.
3. Antonescu C. Round cell sarcomas beyond Ewing: emerging entities. *Histopathology*. 2014 Jan;64(1):26-37.
4. Antonescu CR, Owosho AA, Zhang L, Chen S, Deniz K, Hury JM, Kao YC, Huang SC, Singer S, Tap W, Schaefer IM, Fletcher CD. Sarcomas With CIC-rearrangements Are a Distinct Pathologic Entity With Aggressive Outcome: A Clinicopathologic and Molecular Study of 115 Cases. *Am J Surg Pathol*. 2017 Jul;41(7):941-949.
5. Peters TL, Kumar V, Polikepahad S, Lin FY, Sarabia SF, Liang Y, Wang WL, Lazar AJ, Doddapaneni H, Chao H, Muzny DM, Wheeler DA, Okcu MF, Plon SE, Hicks MJ, López-Terrada D, Parsons DW, Roy A. BCOR-CCNB3 fusions are frequent in undifferentiated sarcomas of male children. *Mod Pathol*. 2015 Apr;28(4):575-86.
6. Peters TL et al. BCOR-CCNB3 fusions are frequent in undifferentiated sarcomas of male children. BCOR-CCNB3 fusions are frequent in undifferentiated sarcomas of male children. *Mod Pathol*. 2015 Apr;28(4):575-86.
7. Cohen-Gogo S et al. Ewing-like sarcomas with BCOR-CCNB3 fusion transcript: A clinical, radiological and pathological retrospective study from the Société Française des Cancers de L'Enfant. *Pediatr Blood Cancer*. 2014 Dec;61(12):2191-8.
8. Puls F et al. BCOR-CCNB3 (Ewing-like) sarcoma: a clinicopathologic analysis of 10 cases, in comparison with conventional Ewing sarcoma. *Am J Surg Pathol*. 2014 Oct;38(10):1307-18.
9. Ludwig K, Alaggio R, Zin A, Peron M, Guzzardo V, Benini S, Righi A, Gambarotti M. BCOR-CCNB3 Undifferentiated Sarcoma-Does Immunohistochemistry Help in the Identification? *Pediatr Dev Pathol*. 2017 Jul-Aug;20(4):321-329.
10. Sugita S, Arai Y, Tonooka A, et al. A novel CIC-FOXO4 gene fusion in undifferentiated small round cell sarcoma: a genetically distinct variant of Ewing-like sarcoma. *Am J Surg Pathol* 2014; 38: 1571-1576.
11. Specht K, Zhang L, Sung YS, et al. Novel BCOR-MAML3 and ZC3H7B-BCOR Gene Fusions in Undifferentiated Small Blue Round Cell Sarcomas. *Am J Surg Pathol* 2016; 40: 433-442.
12. Brohl AS, Solomon DA, Chang W, et al. The genomic landscape of the Ewing Sarcoma family of tumors reveals recurrent STAG2 mutation. *PLoS Genet* 2014; 10: e1004475.
13. Heyer EE, Deveson IW, Wooi D, Selinger CI, Lyons RJ, Hayes VM, O'Toole SA, Ballinger ML, Gill D, Thomas DM, Mercer TR, Blackburn J. Diagnosis of fusion genes using targeted RNA sequencing. *Nat Commun*. 2019 Mar 27;10(1):1388. doi: 10.1038/s41467-019-09374-9. Erratum in: *Nat Commun*. 2020 Apr 8;11(1):1810. PMID: 30918253; PMCID: PMC6437215.